

(12) United States Patent

US 7,601,758 B1 (10) **Patent No.:** Oct. 13, 2009 (45) **Date of Patent:**

(54) METHODS FOR CONCOMITANT ADMINISTRATION OF COLCHICINE AND MACROLIDE ANTIBIOTICS IN THE TREATMENT OF GOUT FLARES

(75) Inventor: Matthew W. Davis, Erwinna, PA (US)

Assignee: Mutual Pharmaceutical Company,

Inc., Philadelphia, PA (US)

Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

(21) Appl. No.: 12/368,700

(22) Filed: Feb. 10, 2009

Related U.S. Application Data

- (63) Continuation of application No. 61/190,053, filed on Oct. 15, 2008.
- (51) **Int. Cl.** A61K 31/167 (2006.01)C07C 233/23 (2006.01)C07C 211/41 (2006.01)
- (52) **U.S. Cl.** **514/629**; 564/123; 564/308; 564/427; 568/306

Field of Classification Search None See application file for complete search history.

References Cited (56)

OTHER PUBLICATIONS

Hung et al., "Fatal Interaction between Clarithromycin and Colchicine in Patients with Renal Insufficiency: A Retrospective Study" Clinical Infectious Diseases (2005) vol. 41 p. 291.*

Brooks, P.M. "Rheumatic Disorders" chapter 25 of Avery's Drug Therapy, pub. 1997 by Adis International, pp. 1113-1161.*

Caraco, et al; "Acute Colchicine Intoxication-Possible Role of Erythromycin Administration"; Journal of Rheumatology; 19; pp. 494-496; (1992).

Horn, et al.; "Life-Threatening Colchicine Drug Interactions"; Pharmacy Times; p. 111; (May 2006).

Perez-Ruiz, et al.; "Optimisation of the Treatment of Acute Gout"; BioDrugs; 13; pp. 415-423; (2000).

Rollot, et al.; "Acute Colchicine Intoxication During Clarithromycin Administration"; The Annals of Pharmacotherapy; 38; pp. 2074-

Terkeltaub, Robert A.; "Gout"; Tne New England Journal of Medicine; 349; pp. 1647-1655; (2003).

van der Velden, et al.; "Colchicine-Induced Neuromyopathy in a Patient with Chronic Renal Failure: The Role of Clarithromycin"; Netherlands Journal of Medicine; 66; pp. 204-206; (2008).

Waknine, Yael; "FDA Safety Changes: Dexedrine and Biaxin CME/ CE"; Medscape Medical News; http://bcbsma.medscape.com/ viewarticle/54411; 5 pages; (2006).

Davis, Matthew W.; Co-pending U.S. Appl. No. 12/327,258; dated Dec. 3, 2009; Title: Methods for Concomitant Administration of Colchicine and Macrolide Antibiotics; Available in the Image File Wrapper (IFW).

Terkeltaub, et al.; "Comparison of Low-Dose vs High-Dose Oral Colchicine Regimens in Patients with Gout Flares"; Presentation at the American College of Rheumatology Scientific Meeting; San Francisco, CA; Oct. 26, 2008.

Terkeltaub, et al.; "The Clinical Relevance of P-gp and CYP 3A4 on Colchicine Metabolism and Elimination"; Presentation at the American College of Rheumatology Scientific Meeting; San Francisco, CA; Oct. 26, 2008.

Krishnan, et al.; "Gout in Ambulatory Care Settings in the United States"; Journal of Rheumatology; 35; pp. 498-501; (2008).

Clarithromycin Tablets, USP; Clarithromycin for Oral Suspension, USP, BIAXIN XL Filmlab; Abbott Laboratories; 07E-010-U780-4 Master; 26 pages; Reference 03-5569-R3 Revised Mar. 2007.

U.S. Appl. No. 12/327,258 Non-final Office Action dated Feb. 13,

U.S. Appl. No. 12/327,258 Final Office Action dated Apr. 16, 2009. U.S. Appl. No. 12/327,258 Non-final Office Action dated Jun. 4,

U.S. Appl. No. 12/327,258 Final Office Action dated Jul. 24, 2009. Morris, et al; "Lesson of the Week-Colchicine in Acute Gout"; BMJ; 327; pp. 1275-1276; (2003).

Leiken et al., "Colchicine"; Poisoning and Toxicology Handbook (4th Ed.); Aug. 2007, p. 216.

"Concomitant Drugs in Medicine"; Concomitant drugs - Wikipedia, http://en.wikipedia.org/wiki/Concomitant_drugs; printed May 28, 2009, 2 pages.

* cited by examiner

Primary Examiner—Shaojia Anna Jiang Assistant Examiner—Eric S Olson (74) Attorney, Agent, or Firm—Cantor Colburn LLP

ABSTRACT

Methods for treating gout flares comprising concomitant administration of colchicine together with one or more macrolide antibiotics, e.g., clarithromycin or erythromycin, are disclosed. Such methods reduce the dangers commonly associated with such concomitant administration and provide additional benefits. Methods of notifying health care practitioners and patients regarding appropriate dosing for co-administration of colchicine together with macrolide antibiotics are also provided.

11 Claims, 2 Drawing Sheets